

Bug Buster-0

- For Use In Organic Gardening
- Provides Rapid Knockdown & Kill of Listed Plant Pests
- Can Be Used on Day of Harvest
- Kills More than 100 Listed Insects



ACTIVE INGREDIENTS:
 Pyrethrins *a botanical insecticide* 1.40%
 OTHER INGREDIENTS: 98.60%
 100.00%

EPA Reg. No. 1021-1771-54705 EPA Est. No. 48498-CA-1

Net Contents: 8 Fl. Oz.
 1 Pint (16 Fl. Oz.)

Distributed by:
 LAWN AND GARDEN PRODUCTS, INC.
 P.O. Box 35000 • Fresno, CA 93745
 (559) 499-2100 • info@montereylawnngarden.com



Contains

PyGanic is a registered trademark of McLaughlin Gormley King.

KEEP OUT OF REACH OF CHILDREN CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration,

	preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For information regarding medical emergencies or pesticide incidents, call 1-888-740-8712.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing vapors or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter storm drains, drainage ditches, gutters or surface waters. Applying this pesticide in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinse application equipment over the treated area will help avoid run off to water bodies or drainage systems. This pesticide is highly toxic to fish. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift from treated areas may be hazardous to organisms in adjacent aquatic sites. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are foraging the treatment area.

TO KILL THE FOLLOWING LISTED INSECTS: including:

12-spotted Cucumber Beetles, Angoumois Grain Moths, Ants (excluding fire and Pharaoh ants), Aphids, Apple Maggots, Armored Scales, Armyworms, Artichoke Plume Moths, Asian Citrus Psyllid, Asparagus Beetles, Bagworms, Bean Beetles, Beet Armyworms, Beetles, Bermudagrass Mirids, Billbugs, Blister Beetles, Blow Flies, Boll Weevils, Box Elder Bugs, Branch and Twig Borers, Cabbage Loopers, Cabbage Maggots, Caddisflies, Cankerworms, Carrot Weevils, Caterpillars, Chalcids, Cheese Skippers, Cherry Fruit Flies, Cigarette Beetles, Clover Mites, Clover Weevils, Cockroaches, Codling Moths, Colorado Potato Beetles, Confused Flour Beetles, Corn Earworms, Crane Flies, Crickets, Cross-striped Cabbageworms, Cucumber Beetles, Dark Mealworms, Darkling Beetles (lesser meal worm), Diamondback Larvae and Moths, Dried Fruit Beetles, Drugstore Beetles, Earwigs, Eastern Tent Caterpillars, Elm Leaf Beetles, Eriophyid Mites, European Pine Tip Moths, Fire worms, Flea Beetles, Forest Tent Caterpillars, Fruit Flies, Fruit tree Leaf rollers, Fruitworms, Fungus Gnats, Garden Symphylan, Glassy Winged Sharpshooters, Gnats, Grain Mites, Grape Leaf Skeletonizers, Grape Leafhoppers, Grape Mealy bugs, Grasshoppers, Green bugs, Green Fruit Worms, Green Peach Aphids, Greenhouse Thrips, Gypsy Moths (adults & larvae), Harlequin Bugs, Heliothis spp., Hornworms, House Flies, Imported Cabbageworms, Indian Meal Moths, Japanese Beetles, Katydid, Lace Bugs, Leaf-footed plant bugs, Leafhoppers, Leafminers, Leafrollers, Leaf-tiers, Loopers, Lygus, Maggots, Mealy Bugs, Mediterranean Flour Moths, Mexican Bean Beetles, Midge, Millipedes, Mites, Mole crickets, Moths, Mushroom Flies, Navel Orangeworms, Olive Fruit Flies, Onion Maggots, Orange Tortrix, Pacific flatheaded borers, Pear Psyllids, Pepper Weevils, Pink bollworms, Potato Leafhoppers, Potato Tuberworms, Proba bugs, Psyllids, Red Flour Beetles, Rice Weevils, Saltmarsh caterpillars, Saw Toothed Grain Beetles, Scales, Sharpshooters, Shore flies, Shot hole borers, Silverfish, Skippers, Small Flying Moths, Soft Scales, Southern chinch bugs, Sow bugs, Spider Beetles, Springtails, Squash bugs, Stable Flies, Stink Bugs, Tarnished Plant Bugs, Thrips, Tobacco Moths, Tomato budworms, Tomato bugs, Tomato Fruit worms, Tomato Hornworms, Tomato pinworms, Tomato Russet Mite, Tussock moths, Vine mealy bugs, Vinegar Flies, Webworms, Weevils, Western Yellow-striped armyworms, Whiteflies, Yellow Mealworms.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS:

- Do not apply this product in a way that will contact workers or other persons, either directly or through drift.
- Do not make applications during the rain.
- Do not wet plants to the point of runoff.
- Not for use in outdoor residential misting systems (indoor or outdoor).
- Do not allow adults, children, or pets to enter the treated area until sprays have dried.
- To avoid possible harm to honey bees, it is advisable to apply in the early morning or late evening hours.
- Not for indoor use.
- Not for use on plants grown for sale.

Pyrethrins degrade rapidly in sunlight. To ensure the best performance, coverage is key. Use spray equipment that will provide the best coverage and direct contact with as many insects as possible.

Dilute 1 to 1.4 fl. oz. of Bug Buster-O per gallon of water per 1,000 sq. ft. Do not exceed the maximum application rates of 1.4 fl. oz. per 1,000 sq. ft. Apply dilute solution with all types of agricultural sprayers, including compressed air sprayers or pump driven sprayers. Mix only enough for immediate use. Spraying should begin when the insects first appear. Do not wait until the plants are heavily infested.

IMPORTANT NOTE: Plant safety is an important consideration when using insecticides. However, it is not possible to evaluate the phytotoxicity of Bug Buster-O on numerous plant varieties that may react differently to insecticides in different growth stages or under varying environmental conditions. Before making widespread applications of Bug Buster-O, treat a limited number of plants and observe for phytotoxicity over a 10 day period.

ROOT AND TUBER VEGETABLES: Including: Arracacha, Arrowroot, Purple Arrowroot, Japanese Artichoke, Jerusalem Artichoke, Garden Beets, Sugar Beets, Edible Burdock, Edible Canna, Carrots, Cassava (bitter or sweet), Celeriac (celery root), Chayote (root), Chervil (turnip rooted), Chicory, Chufa, Dasheen, Ginger, Ginseng, Horseradish, Leren, Parsley (turnip rooted), Parsnip, Potato, Radish, Japanese Radish (Daikon), Rutabaga, Salsify, Black Salsify, Spanish Salsify, Sweet Potato, Tanier, Turmeric, Turnip, Yam (true), Yam Bean.

LEAVES OF ROOT AND TUBER VEGETABLES: Including: Garden Beet, Sugar Beet, Edible Burdock, Carrot, Cassava (bitter or sweet), Celeriac (celery root), Chervil (Turnip Rooted), Chicory, Dasheen (Taro), Parsnip, Radish, Japanese Radish (Daikon), Rutabaga, Black Salsify, Sweet Potato, Tanier, Turnip, Yam (true).

BULB VEGETABLES: Including: Garlic, Great-headed Garlic, Leek, Onion (bulb and green), Welc h, Shallot.

LEAFY VEGETABLES: Including: Amaranth (Leafy Amaranth, Chinese Spinach, Tampala), Arugula, Cardoon, Celery, Chinese Celery, Celtuce, Chervil, Cilantro, Corn Salad, Chrysanthemum (edible-leaved), Chrysanthemum (garland), Cress (garden, water), Upland Cress (yellow rocket, winter cress), Dandelion, Dock (sorrel), Endive (escarole), Fennel (Florence), Lettuce (head and leafy), Orach, Parsley, Purslane (garden & winter), Radicchio, Rhubarb, Spinach, Fine Spinach (Malabar, Ceylon), Spinach (New Zealand), Swiss Chard.

BRASSICA (COLE) LEAFY VEGETABLES: Including: Broccoli, Chinese Broccoli (Gai Lon), Broccoli raab (Rapini), Brussels Sprouts, Cabbage, Chinese Cabbage (Bok Choy), Chinese Cabbage (Napa), Chinese Mustard Cabbage (Gai Choy), Cauliflower, Cavallo broccolo, Collards, Kale, Kohlrabi, Mizuna, Mustard Greens, Mustard Spinach, Rape Greens.

LEGUME VEGETABLES (SUCCULENT OR DRIED): Including: Adzuki Beans, Field Beans, Kidney Beans, Lima Beans, Moth Beans, Mung Beans, Navy Beans, Pinto Beans, Rice Beans, Runner Beans, Snap Beans, Tepary Beans, Urd Beans, Wax Beans, Asparagus Beans, Black-eyed Peas, Catjang, Chinese Longbeans, Cowpeas, Crowder Peas, Southern Peas, Yard-Longbeans, Broad Beans, (Fava Beans), Chick Peas (Garbanzo Beans), Guar, Jack Bean (Sword Bean), Lablab Bean (Hyacinth Bean), Lentils, Peas (Garden Peas), Field Peas, Sugar

Snap Peas, English Pea, Snow Pea), Pigeon Peas, Soybeans, Sweet Lupin Beans, White Lupin Beans, White Sweet Lupin, Sword Bean.

FOLIAGE OF LEGUME VEGETABLES: Including: Plant part of any legume vegetable included in the legume vegetable group that will be used as animal feed, including any variety of Beans, Field Peas, Soybeans.

FRUITING VEGETABLES: Including: Eggplant, Ground Cherry, Okra, Pepinos, Pepper (Bell Pepper, Chili Pepper, Cooking Peppers, Pimentos, Sweet Peppers), Tomatillo, Tomatoes.

CUCURBIT VEGETABLES: Including: Balsam Apple, Balsam Pear (Bitter Melon), Chayote (fruit), Chinese Waxgourd (Chinese preserving melon), Chinese Cucumber, Citron Melon, Cucumber, Gherkin, Edible Gourds, Melons (including hybrids, Cantaloupe, Casaba, Crenshaw, Golden Pershaw Melon, Honeydew Melons, Honey Balls, Mango Melon, Muskmelon, Persian Melon, Pineapple Melon, Santa Claus Melon, Snake Melon), Pumpkin, Squash (summer & winter), Watermelon (including hybrids).

CITRUS FRUITS: Including: Calamondin, Citrus Citron, Citrus Hybrids, Grapefruit, Kumquats, Lemons, Limes, Mandarin (Tangerine), Orange (sweet & sour), Pummelo, Satsuma Mandarin, (Citrus spp. Includes Chironja, Tangelos, Tangors).

POME FRUITS: Including: Apple, Crabapple, Loquat, Pear, Mayhaw, Oriental Pear, Quince.

STONE FRUITS: Including: Apricot, Cherry (sweet & sour), Nectarine, Peach, Plum, Prune, Chickasaw Plum, Damson Plum, Japanese Plum, Plumcot.

SMALL FRUITS AND BERRIES: Including: Blackberry, Blueberry, Cranberry, Currant, Dewberry, Elderberry, Gooseberry, Grape, Huckleberry, Loganberry, Olallieberry, Raspberry (black & red), Strawberry, Youngberry.

TREE NUTS: Including: Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (hazelnut), Hickory nut, Macadamia Nut (Bush Nut), Pecan, Pistachio, Walnut, Black and English (Persian).

ORIENTAL VEGETABLES: Including: Acerola, Atemoya, Balsam Pear (bitter melon), Carambola, Japanese Artichoke, Chinese Broccoli (Gai Lan), Chinese Cabbage (Bok Choy, Napa), Chinese Mustard Cabbage (Gai Choy), Dasheen, Ginger, Ginseng, Chinese Longbeans, Mung Beans, Citron Melon, Japanese Radish (Daikon), Chinese Spinach, Chinese Waxgourd, Cilantro, Citron Melon, Rambutan, Water Chestnut.

SUBTROPICAL FRUITS: Including: Avocado, Banana, Carob, Barbados Cherry, Cherimoya, Dates, Durian (Jackfruit), Feijoa, Figs, Guava, Kiwifruit, Lychee, Mango, Papaya, Passion Fruit, Persimmon, Pineapple, Pomegranate.

HERBS AND SPICES: Including: Allspice, Angelica, Anise (anise seed), Anise [star], Annatto (seed), Balm (lemon balm), Basil, Borage, Burnet, Camomile, Capers buds, Caraway, Caraway [black], Cardamom, Cassia bark, Cassia buds, Catnip, Celery seed, Chervil (dried), Chicory, Chive, Chive [Chinese], Cinnamon, Clary, Clove buds, Coriander (cilantro or Chinese parsley) (leaf), Coriander (cilantro) (seed), Costmary, Culantro (leaf), Culantro (seed), Cumin, Curry (leaf), Dill (dill weed), Dill (seed), Fennel (common), Fennel [Florence] (seed), Fenugreek, Grains of paradise, Horehound, Hyssop, Juniper berry, Lavender, Lemongrass, Lovage (leaf), Lovage (seed), Mace, Marigold, Marjoram (includes sweet or annual marjoram, wild marjoram or oregano, and pot marjoram), Mustard (seed), Nasturtium, Nutmeg, Oregano, Mint, Paprika, Parsley (dried), Pennyroyal, Pepper [black], Pepper [white], Poppy (seed), Rosemary, Rue, Saffron, Sage, Savory [summer and winter], Sweet bay (bay leaf), Tansy, Tarragon, Thyme, Vanilla, Wintergreen, Woodruff, Wormwood.

ADDITIONAL PLANTS: Including: Artichoke, Asparagus, Avocado, Coffee, Cotton, Hops, Jojoba, Mushroom, Okra, Olives, Peanuts, Pineapple, Rice, Safflowers, Sesame, Sugar Cane, Sunflower, Tea.

FOR USE ON ORNAMENTAL PLANTS GROWN IN GARDENS: Including: African Violet, Ageratum, Aster, Azalea, Begonia, Calceolaria,

Calendula, Calla, Camellia, Carnation, Ceanothus, Cineraria, Chrysanthemum, Coleus, Cyclamen, Cypress, Daffodil, Dahlia, Delphinium, Dogwood, Elm, Eucalyptus, Fern, Ficus, Foliage Plants, Fuchsia, Gardenia, Geranium, Gladiolus, Gloxinia, Gypsophila, Holly, Hyacinth, Hydrangea, Iris, Lily, Maidenhair Fern, Marigold, Narcissus, Palm, Pansy, Peony, Pelargonium, Petunia, Philodendron, Phlox, Pine, Pyracantha, Rhododendron, Roses, Rubber Plant, Snapdragon, Sweet Pea, Tulips, Viburnum, Wandering Jew, Yew, Zinnia and Andromeda, Arborvitae, Ash, Beech, Birch, Boxwood, Cotoneaster, Crabapple, Euonymus, Fir, Firethorn, Forsythia, Hawthorn, Hemlock, Hickory, Honey Locust, Horse Chestnut, Juniper, Larch, Laurel, Lilac, Linden, Mimosa, Myrtle, Oak, Pine, Privet, Tulip Tree, Viburnum, Willow.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Keep this product in its tightly closed original container when not in use. Store in a cool, dry (preferably locked) area that is inaccessible to children and animals.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Nonrefillable container. Do not reuse or refill this container. **If empty:** offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

(MGK-032816N) 0913/0316(02)